




ELSEVIER

Molecular and Cellular Endocrinology 108 (1995) 75–85



## Production of a biologically active recombinant teleostean growth hormone in *E. coli* cells

Clara M. Cheng<sup>a,b</sup>, C.M. Lin<sup>a</sup>, M. Shablott<sup>a,b</sup>, L.I. Gonzalez-Villasenor<sup>c</sup>, D.A. Powers<sup>d</sup>, C. Woods<sup>e</sup>,  
T.T. Chen<sup>a,b,\*</sup>

<sup>a</sup>Center of Marine Biotechnology, University of Maryland Biotechnology Institute, 600 E. Lombard Street, Baltimore, MD 21202, USA

<sup>b</sup>Department of Biological Sciences, University of Maryland at Baltimore County, Baltimore, MD 21228, USA

<sup>c</sup>Department of Biology, The Johns Hopkins University, 34th and N. Charles Streets, Baltimore, MD 21218, USA

<sup>d</sup>Hopkins Marine Station, Stanford University, Pacific Grove, CA 93590, USA

<sup>e</sup>Agriculture Experimental Station, University of Maryland System, College Park, MD, USA

Received 30 September 1994; accepted 5 December 1994

### Abstract

We have isolated and characterized several recombinant lambda phage clones carrying growth hormone (GH) cDNA of striped bass (*Morone saxatilis*). Nucleotide sequence and the predicted amino acid sequence of sbGH was determined from a recombinant clone carrying the longest cDNA insert. The sbGH cDNA encodes a pre-hormone of 204 amino acid residues. Comparison of the predicted amino acid sequence of sbGH with those of other vertebrates revealed different degrees of sequence identity: approximately 98% with European sea bass; 90% with bluefin tuna; bonito and red seabream; 71% with winter flounder; 64% with salmonids; 55% with carp; and 38% with human. Expression of the mature sbGH cDNA (without the signal peptide sequence) in *E. coli* cells under regulation of the lambda phage PL promoter produced a polypeptide of 20 kDa. Following renaturation, this recombinant hormone was shown to be biologically active in a radioreceptor competition binding assay and in the induction of hepatic insulin-like growth factor I (IGF-I) mRNA synthesis in vivo.

**Keywords:** Striped bass; Growth hormone; Expression in *E. coli*; Renaturation; Competitive receptor binding assay; IGF-I induction